


CLAIMS


1. A quinazoline derivative of the Formula I



- 5 wherein Z is an O, S, SO, SO₂, N(R²) or C(R²)₂ group, wherein each R² group, which may be the same or different, is hydrogen or (1-6C)alkyl;

m is 0, 1, 2, 3 or 4;

- each R¹ group, which may be the same or different, is selected from halogeno, trifluoromethyl, cyano, isocyano, nitro, hydroxy, mercapto, amino, formyl, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, (3-6C)alkenoylamino, N-(1-6C)alkyl-(3-6C)alkenoylamino, (3-6C)alkynoylamino, N-(1-6C)alkyl-(3-6C)alkynoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino, N-(1-6C)alkyl-(1-6C)alkanesulphonylamino or from a group of the formula :




- wherein X¹ is a direct bond or is selected from O, S, SO, SO₂, N(R⁴), CO, CH(OR⁴), CON(R⁴), N(R⁴)CO, SO₂N(R⁴), N(R⁴)SO₂, OC(R⁴)₂, SC(R⁴)₂ and N(R⁴)C(R⁴)₂, wherein R⁴ is hydrogen or (1-6C)alkyl, and Q¹ is aryl, aryl-(1-6C)alkyl, (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, (3-7C)cycloalkenyl, (3-7C)cycloalkenyl-(1-6C)alkyl, heteroaryl, heteroaryl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl, or (R¹)_m is (1-3C)alkylenedioxy,

- and wherein adjacent carbon atoms in any (2-6C)alkylene chain within a R¹ substituent are optionally separated by the insertion into the chain of a group selected from O, S, SO, SO₂, N(R⁵), CO, CH(OR⁵), CON(R⁵), N(R⁵)CO, SO₂N(R⁵), N(R⁵)SO₂, CH=CH and C≡C wherein

- 153 -


R⁵ is hydrogen or (1-6C)alkyl or, when the inserted group is N(R⁵), R⁵ may also be (2-6C)alkanoyl,

and wherein any CH₂=CH- or HC≡C- group within a R¹ substituent optionally bears at the terminal CH₂= or HC≡ position a substituent selected from halogeno, carboxy, carbamoyl, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl or from a group of the formula :



wherein X² is a direct bond or is selected from CO and N(R⁶)CO, wherein R⁶ is hydrogen or (1-6C)alkyl, and Q² is aryl, aryl-(1-6C)alkyl, heteroaryl, heteroaryl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl,

and wherein any CH₂ or CH₃ group within a R¹ substituent optionally bears on each said CH₂ or CH₃ group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, amino, carboxy, carbamoyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxyalkylamino di-[(1-6C)alkoxyalkyl]amino or hydroxy(1-6C)alkylamino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino, N-(1-6C)alkyl-(1-6C)alkanesulphonylamino or from a group of the formula :




wherein X³ is a direct bond or is selected from O, S, SO, SO₂, N(R⁷), CO, CH(OR⁷), CON(R⁷), N(R⁷)CO, SO₂N(R⁷), N(R⁷)SO₂, C(R⁷)₂O, C(R⁷)₂S and N(R⁷)C(R⁷)₂, wherein R⁷ is hydrogen or (1-6C)alkyl, and Q³ is aryl, aryl-(1-6C)alkyl, (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, (3-7C)cycloalkenyl, (3-7C)cycloalkenyl-(1-6C)alkyl, heteroaryl, heteroaryl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl,

and wherein any aryl, heteroaryl or heterocyclyl group within a substituent on R¹ optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, formyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino,


- 154 -

di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl,
N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino,
N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl,
N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino, N-(1-6C)alkyl-

5 (1-6C)alkanesulphonylamino or from a group of the formula :



wherein X^4 is a direct bond or is selected from O and N(R^9), wherein R^9 is hydrogen or
 (1-6C)alkyl, and R^8 is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl,
 cyano-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-
 10 (1-6C)alkyl, (2-6C)alkanoylamino-(1-6C)alkyl, (1-6C)alkoxycarbonylamino-(1-6C)alkyl,
 or a group of the formula :



wherein X^5 is a direct bond or is selected from O, N(R^{10}) and CO, wherein R^{10} is hydrogen or
 (1-6C)alkyl, and Q^4 is aryl, aryl-(1-6C)alkyl, heteroaryl, heteroaryl-(1-6C)alkyl, heterocyclyl
 15 or heterocyclyl-(1-6C)alkyl which optionally bears 1 or 2 substituents, which may be the same
 or different, selected from halogeno, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl and
 (1-6C)alkoxy,

and wherein any heterocyclyl group within a substituent on R^1 optionally bears 1 or 2
 oxo or thioxo substituents;

20 n is 0, 1 or 2; and

R^3 is selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy,
 carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-
 6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl,
 (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl,
 25 N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino,
N-(1-6C)alkyl-(2-6C)alkanoylamino, (3-6C)alkenoylamino, N-(1-6C)alkyl-
 (3-6C)alkenoylamino, (3-6C)alkynoylamino, N-(1-6C)alkyl-(3-6C)alkynoylamino,
N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino, N-
 (1-6C)alkyl-(1-6C)alkanesulphonylamino or from a group of the formula :

30 $-X^6-R^{11}$


wherein X^6 is a direct bond or is selected from O and N(R^{12}), wherein R^{12} is hydrogen or
 (1-6C)alkyl, and R^{11} is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl,

- 155 -


cyano-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl or di-[(1-6C)alkyl]amino-(1-6C)alkyl;

Z^2 is a $C\equiv C$ or $C(R^{13})=C(R^{13})$ group, wherein each R^{13} group, which may be the same or different, is hydrogen or (1-6C)alkyl; and

- 5 R^{14} is selected from halogeno, cyano, isocyano, formyl, carboxy, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-
10 (1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, (2-6C)alkanoylamino-(1-6C)alkyl, (1-6C)alkoxycarbonylamino-(1-6C)alkyl, from a group of formula:




- wherein x is 0, 1, 2 or 3, R^c is hydrogen or (1-6C)alkyl and R^a and R^b are each independently selected from hydrogen and (1-6C)alkyl or R^a and R^b together with the nitrogen to which they
15 are attached form a 4 to 7 membered heterocyclyl optionally containing up to two further heteratoms selected from oxygen, nitrogen or sulphur, or from a group of the formula :



- wherein X^7 is a direct bond or is selected from CO , $CH(OR^{15})$, $CON(R^{15})$ or $SO_2N(R^{15})$, wherein R^{15} is hydrogen or (1-6C)alkyl, and Q^5 is aryl, aryl-(1-6C)alkyl, (3-7C)cycloalkyl,
20 (3-7C)cycloalkyl-(1-6C)alkyl, heteroaryl, heteroaryl-(1-6C)alkyl, heterocyclyl heterocyclyl-(1-6C)alkyl or heterocyclyloxy-(1-6C)alkyl,

- and wherein any CH , CH_2 or CH_3 group within a R^{14} substituent optionally bears on each said CH , CH_2 or CH_3 group one or more halogeno, (1-6C)alkyl or (3-6C)cycloalkyl substituents or a substituent selected from hydroxy, cyano, amino, carboxy, carbamoyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino,
25 di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino,
30 N-(1-6C)alkyl-(1-6C)alkanesulphonylamino or from a group of the formula :



- 156 -


wherein X^8 is a direct bond or is selected from O, S, SO, SO_2 , $N(R^{16})$, CO, $CH(OR^{16})$, $CON(R^{16})$, $N(R^{16})CO$, $SO_2N(R^{16})$, $N(R^{16})SO_2$, $C(R^{16})_2O$, $C(R^{16})_2S$ and $N(R^{16})C(R^{16})_2$, wherein R^{16} is hydrogen or (1-6C)alkyl, and Q^6 is aryl, aryl-(1-6C)alkyl, (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, (3-7C)cycloalkenyl, (3-7C)cycloalkenyl-(1-6C)alkyl,

5 heteroaryl, heteroaryl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl,

and wherein any aryl, heteroaryl or heterocyclyl group within a substituent on R^{14} optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy,


10 (1-6C)alkylthio, (1-6C)alkylsulphinyl, (1-6C)alkylsulphonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulphamoyl, N,N-di-[(1-6C)alkyl]sulphamoyl, (1-6C)alkanesulphonylamino, N-(1-6C)alkyl-

15 (1-6C)alkanesulphonylamino or from a group of the formula :



wherein X^9 is a direct bond or is selected from O and $N(R^{18})$, wherein R^{18} is hydrogen or (1-6C)alkyl, and R^{17} is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-

20 (1-6C)alkyl, (2-6C)alkanoylamino-(1-6C)alkyl, (1-6C)alkoxycarbonylamino-(1-6C)alkyl, or from a group of the formula :



wherein X^{10} is a direct bond or is selected from O, $N(R^{19})$ and CO, wherein R^{19} is hydrogen or (1-6C)alkyl, and Q^7 is aryl, aryl-(1-6C)alkyl, heteroaryl, heteroaryl-(1-6C)alkyl, heterocyclyl

25 or heterocyclyl-(1-6C)alkyl which optionally bears 1 or 2 substituents, which may be the same or different, selected from halogeno, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl and (1-6C)alkoxy,


and wherein any heterocyclyl group within a substituent on R^{14} optionally bears 1 or 2 oxo or thioxo substituents;

30 or a pharmaceutically-acceptable salt thereof.

- 157 -

2. A quinazoline derivative of the Formula I or a pharmaceutically acceptable salt thereof, according to claim 1 wherein R^1 , R^2 , R^3 , Z , Z^2 , m and n have any of the meanings defined in claim 1 and

R^{14} is selected from hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, from a group of formula:




wherein x is 0, 1, 2 or 3, R^c is hydrogen or (1-6C)alkyl and R^a and R^b are each independently selected from hydrogen and (1-6C)alkyl or R^a and R^b together with the nitrogen to which they are attached form morpholino, or from a group of the formula :



wherein X^7 is a direct bond and Q^5 is aryl, heteroaryl, heterocyclyl, heterocyclyl-(1-6C)alkyl or heterocycloxy-(1-6C)alkyl,

and wherein any CH , CH_2 or CH_3 group within a R^{14} substituent optionally bears on each said CH , CH_2 or CH_3 group one or more halogeno, (1-6C)alkyl, (1-6C)alkoxy, carbamoyl, (1-6C)alkoxy, \underline{N} -(1-6C)alkylcarbamoyl, \underline{N} , \underline{N} -di-[(1-6C)alkyl]carbamoyl or from a group of the formula :



wherein X^8 is a direct bond or O and Q^6 is aryl, (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl,


and wherein any aryl, cycloalkyl or heterocyclyl group within a substituent on R^{14} optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from halogeno, hydroxyl, amino, (1-6C)alkyl or (1-6C)alkanoyl,

and wherein any heterocyclyl group within a substituent on R^{14} optionally bears 1 or 2 oxo or thioxo substituents.

25

3. A quinazoline derivative of the Formula I according to claim 1 wherein Z is O or NH

m is 1 and the R^1 group is located at the 5-, 6-, or 7-position or m is 2 and each R^1 group, which may be the same or different, is located at the 5- and 7-positions or at the 6- and 7-positions and R^1 is selected from methoxy, ethoxy, propoxy, isopropoxy, 2-methylpropoxy, or from a group of the formula :



- 158 -

wherein X^1 is O and Q^1 is piperidino, piperidin-3-yl, piperidin-4-yl, 1-, 3- or 4-homopiperidinyl, piperazin-1-yl, homopiperazin-1-yl, 1-, 2- or 3-pyrrolidinylmethyl, morpholinomethyl, piperidinomethyl, 3- or 4-piperidinylmethyl, 1-, 3- or 4-homopiperidinylmethyl, 2-pyrrolidin-1-ylethyl, 3-pyrrolidin-2-ylpropyl, pyrrolidin-2-ylmethyl,
 5 2-pyrrolidin-2-ylethyl, 3-pyrrolidin-1-ylpropyl, 4-pyrrolidin-1-ylbutyl, 2-morpholinoethyl, 3-morpholinopropyl, 4-morpholinobutyl, tetrahydro-2H-pyran-4-yl, 2-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)ethyl, 3-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)propyl, 2-piperidinoethyl, 3-piperidinopropyl, 4-piperidinobutyl, piperidin-4-ylmethyl, 2-piperidin-3-ylethyl, 3-piperidin-1-ylpropyl, 3-piperidin-3-ylpropyl, 2-piperidin-4-ylethyl, 3-piperidin-4-ylpropyl, 2-
 10 homopiperidin-1-ylethyl, 3-homopiperidin-1-ylpropyl, 2-piperazin-1-ylethyl, 3-piperazin-1-ylpropyl, 4-piperazin-1-ylbutyl, 2-homopiperazin-1-ylethyl or 3-homopiperazin-1-ylpropyl,

and wherein adjacent carbon atoms in any (2-6C)alkylene chain within a R^1 substituent are optionally separated by the insertion into the chain of a O,

and wherein any CH_2 or CH_3 group within a R^1 substituent optionally bears on each
 15 said CH_2 or CH_3 group one or more fluoro, chloro or bromo groups or a substituent selected from amino, methylamino, dimethylamino, methoxyethylamino, di-(methoxyethyl)amino, or hydroxypropylamino,


and wherein any heterocyclyl group within a substituent on R^1 optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from methyl, ethyl, methoxy,
 20 ethoxy, formyl, acetyl, hydroxyl, hydroxymethyl, fluoroethyl or hydroxypropylamino,

and wherein any heterocyclyl group within a substituent on R^1 optionally bears 1 or 2 oxo substituents;

n is 0 or 1 and R^3 group, if present, is located at the 5- or 6-position of the 1,3-benzodioxol-4-yl group and is selected from fluoro, chloro, bromo, trifluoromethyl, cyano,
 25 hydroxy, methyl, ethyl, methoxy and ethoxy;


Z^2 is a $C\equiv C$ or $CH=CH$ group; and

R^{14} is selected from hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, from a group of formula:




30 wherein x is 0, 1, 2 or 3, R^c is hydrogen or (1-6C)alkyl and R^a and R^b are each independently selected from hydrogen and (1-6C)alkyl or R^a and R^b together with the nitrogen to which they are attached form morpholino or from a group of the formula :

- 159 -



wherein X^7 is a direct bond and Q^5 is aryl, heteroaryl, heterocyclyl, heterocyclyl-(1-6C)alkyl or heterocycliloxy-(1-6C)alkyl,

and wherein any CH, CH₂ or CH₃ group within a R^{14} substituent optionally bears on
 5 each said CH, CH₂ or CH₃ group one or more halogeno, (1-6C)alkyl, (1-6C)alkoxy, carbamoyl, (1-6C)alkoxy, N-(1-6C)alkylcarbamoyl, N, N-di-[(1-6C)alkyl]carbamoyl or from a group of the formula :



wherein X^8 is a direct bond or O and Q^6 is aryl, (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-
 10 6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl,

and wherein any aryl, cycloalkyl or heterocyclyl group within a substituent on R^{14} optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from halogeno, hydroxyl, amino, (1-6C)alkyl or (1-6C)alkanoyl,

and wherein any heterocyclyl group within a substituent on R^{14} optionally bears 1 or 2
 15 oxo or thioxo substituents;

or a pharmaceutically acceptable acid addition salt thereof.

4. A quinazoline derivative of the Formula I, or a pharmaceutically acceptable salt thereof, according to claim 1 wherein R^1 , R^2 , R^3 , R^{14} , Z^2 , m and n have any of the meanings
 20 defined in claim 1 and Z is NH.

5. A quinazoline derivative of the Formula I according to claim 1 wherein Z is NH

m is 2,

25 and the first R^1 group is a 6-methoxy group and the second R^1 group is located at the 7-position and is selected from methoxy, 2-methoxyethoxy, 2-fluoroethoxy, 3-chloroethoxy, isopropoxy, isopropylmethoxy, 3-dimethylaminopropoxy, 2-dimethylaminoethoxy, dimethylaminoisopropoxy, 2-[3-(hydroxy)propylamino]ethoxy, 3-[bis(2-methoxyethyl)amino]propoxy, 2-(2-methoxyethoxy)ethoxy, 3-chloropropoxy, 2-(2-
 30 chloroethoxy)ethoxy, 3-piperazin-1-ylpropoxy, 2-piperazin-1-ylethoxy, 4-piperazin-1-ylbutoxy, 2-(4-methylpiperazin-1-yl)ethoxy, 3-(4-methylpiperazin-1-yl)propoxy, 4-(4-methylpiperazin-1-yl)butoxy, 3-[4-(2-fluoroethyl)piperazin-1-yl]propoxy, 2-[4-(2-

- 160 -

fluoroethyl)piperazin-1-yl]ethoxy, 4-[4-(2-fluoroethyl)piperazin-1-yl]butoxy, 3-(4-acetylpiperazin-1-yl)propoxy, 2-(4-acetylpiperazin-1-yl)ethoxy, 4-(4-acetylpiperazin-1-yl)butoxy, 3-(4-formylpiperazin-1-yl)propoxy, 2-(4-formylpiperazin-1-yl)ethoxy, 4-(4-formylpiperazin-1-yl)butoxy, 3-morpholinopropoxy, 2-morpholin-4-ylethoxy, 4-morpholin-4-ylbutoxy, 3-(2,6-dimethylmorpholin-4-yl)propoxy, 2-(2,6-dimethylmorpholin-4-yl)ethoxy, 4-(2,6-dimethylmorpholin-4-yl)butoxy, 3-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)propoxy, 3-[2-(hydroxymethyl)pyrrolidin-1-yl]propoxy, 4-[2-(hydroxymethyl)pyrrolidin-1-yl]butoxy, 2-[2-(hydroxymethyl)pyrrolidin-1-yl]ethoxy, 3-(4-hydroxypiperidin-1-yl)propoxy, 2-(4-hydroxypiperidin-1-yl)ethoxy, 4-(4-hydroxypiperidin-1-yl)butoxy, 1-methylpiperidin-4-ylmethoxy, 3-(1-methylpiperidin-4-yl)propoxy, 3-(4-methoxypiperidin-1-yl)propoxy, 3-(4-methoxypiperidin-1-yl)ethoxy or 4-(4-methoxypiperidin-1-yl)butoxy

and wherein any heterocyclyl group within a substituent on R¹ optionally bears 1 or 2 oxo substituents;


n is 0 or 1 and R³ group, if present, is located at the 5-position of the 1,3-benzodioxol-yl group and is selected from fluoro, chloro, bromo, trifluoromethyl, cyano, hydroxy, methyl, ethyl, methoxy and ethoxy;

Z² is a C≡C or CH=CH group; and

R¹⁴ is selected from (1-6C)alkoxy-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, hydroxyl-(1-6C)alkyl or from a group of formula:



wherein x is 0, 1, 2 or 3, R^c is hydrogen or (1-6C)alkyl and R^a and R^b are each independently selected from hydrogen and (1-6C)alkyl or R^a and R^b together with the nitrogen to which they are attached form morpholino, or from a group of the formula :



wherein X⁷ is a direct bond and Q⁵ is aryl, heteroaryl, heterocyclyl, heterocyclyl-(1-6C)alkyl, or heterocycliloxy-(1-6C)alkyl,

and wherein any CH, CH₂ or CH₃ group within a R¹⁴ substituent optionally bears on each said CH, CH₂ or CH₃ group a substituent selected from (1-6C)alkoxy, carbamoyl, N-(1-6C)alkylcarbamoyl or a group of the formula :




wherein X⁸ is a direct bond or O and Q⁶ is (3-7C)cycloalkyl or heterocyclyl

- 161 -

and wherein any heterocyclyl group within a substituent on R¹⁴ optionally bears 1 oxo substituent;

or a pharmaceutically acceptable acid addition salt thereof.

- 5 6. A quinazoline derivative of the Formula I, or a pharmaceutically acceptable acid addition salt thereof, according to claim 1 wherein R¹, R², R³, Z, Z², m and n have any of the meanings defined in claim 1 and R¹⁴ is a group of the formula:



wherein x is 1, R^c is hydrogen or (1-3C) alkyl and R^a and R^b are each independently selected
10 from hydrogen and (1-3C)alkyl.

7. A quinazoline derivative of the Formula I according to claim 1 wherein Z is NH

m is 2,

- 15 and the first R¹ group is a 6-methoxy group and the second R¹ group is located at the 7-position and is selected from methoxy, 2-methoxyethoxy, isopropoxy, isopropylmethoxy, 3-dimethylaminopropoxy, dimethylaminoisopropoxy, 2-[3-(hydroxy)propylamino]ethoxy, 3-[bis(2-methoxyethyl)amino]propoxy, 2-(4-methylpiperazin-1-yl)ethoxy, 3-(3-oxo-4-methylpiperazin-1-yl)propoxy, 3-(2-oxo-4-methylpiperazin-1-yl)propoxy, 3-morpholinopropoxy, 2-morpholin-4-ylethoxy, 4-morpholin-4-ylbutoxy, 3-(2,6-dimethylmorpholin-4-yl)propoxy, 3-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)propoxy, 2-fluoroethoxy, 3-[4-(2-fluoroethyl)piperazin-1-yl]propoxy, 3-(4-acetylpiperazin-1-yl)propoxy, 3-(4-formylpiperazin-1-yl)propoxy, 3-piperazin-1-ylpropoxy, 3-(4-hydroxypiperidin-1-yl)propoxy, 3-[2-(hydroxymethyl)pyrrolidin-1-yl]propoxy, 2-(2-methoxyethoxy)ethoxy, 3-chloropropoxy, 2-(2-chloroethoxy)ethoxy, 1-methylpiperidin-4-ylmethoxy, 3-(4-methoxypiperidin-1-yl)propoxy or 3-(4-hydroxypiperidin-1-yl)propoxy,


n is 1 and R³ group, if present, is located at the 6 position of the 1,3-benzodioxol group and is selected from fluoro, chloro or bromo;

Z² is a C≡C or CH=CH group; and

- 30 R¹⁴ is selected from methoxymethyl, 1-methoxyethyl, 2-methoxyethyl, methoxyisopropyl, 2-methoxypropyl, ethoxymethyl, methoxyethoxymethyl, hydroxymethyl, carbamoylmethoxymethyl, methylcarbamoylmethoxymethyl, isopropoxymethyl, di-

- 162 -


(methylamino)methyl, hydroxyisopropyl, (cyclopropylmethoxy)methyl, (cyclopentylmethoxy)methyl from a group of formula:



- wherein x is 1, R^c is hydrogen and R^a and R^b are each independently selected from hydrogen, and methyl or R^a and R^b together with the nitrogen to which they are attached form morpholino, or x is 1 and R^a, R^b and R^c are all methyl, or is selected from 2-oxo-pyrrolidin-1-ylmethyl, pyridin-2-yl, (tetrahydrofuran-3-ylmethoxy)methyl, (tetrahydrofuran-3-yloxy)methyl, [(1,3-dioxolan-2-yl)methoxy]methyl, phenyl, pyridin-3-yl, pyrazin-3-yl, pyrimidin-2-yl, 1H-pyrazol-4-yl or 1H-pyrazol-5-yl;
- or a pharmaceutically acceptable acid addition salt thereof.

8. A quinazoline derivative of the Formula I according to claim 1 wherein Z is NH

- m is 2 and each R¹ group, which may be the same or different, is located at the 5- and 7-positions and R¹ is selected from methoxy, ethoxy, propoxy, isopropoxy, 2-methylpropoxy, vinyloxy, or from a group of the formula :



- wherein X¹ is O and Q¹ is 1-, 2-, or 3-pyrrolidinyl, piperidino, piperidin-3-yl, piperidin-4-yl, 1-, 3- or 4-homopiperidinyl, piperazin-1-yl, homopiperazin-1-yl, 1-, 2- or 3-pyrrolidinylmethyl, morpholinomethyl, piperidinomethyl, 3- or 4-piperidinylmethyl, 1-, 3- or 4-homopiperidinylmethyl, 2-pyrrolidin-1-ylethyl, 3-pyrrolidin-2-ylpropyl, pyrrolidin-2-ylmethyl, 2-pyrrolidin-2-ylethyl, 3-pyrrolidin-1-ylpropyl, 4-pyrrolidin-1-ylbutyl, 2-morpholinoethyl, 3-morpholinopropyl, 4-morpholinobutyl, tetrahydro-2H-pyran-4-yl, 2-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)ethyl, 3-(1,1-dioxotetrahydro-4H-1,4-thiazin-4-yl)propyl, 2-piperidin-2-ylethyl, 3-piperidin-1-ylpropyl, 3-piperidin-3-ylpropyl, 2-piperidin-4-ylethyl, 3-piperidin-4-ylpropyl, 2-homopiperidin-1-ylethyl, 3-homopiperidin-1-ylpropyl, 2-piperazin-1-ylethyl, 3-piperazin-1-ylpropyl, 4-piperazin-1-ylbutyl, 2-homopiperazin-1-ylethyl or 3-homopiperazin-1-ylpropyl,
- and wherein any CH₂ or CH₃ group within a R¹ substituent optionally bears on each said CH₂ or CH₃ group one or more fluoro or chloro groups or a substituent selected from

- 163 -

amino, methylamino, methoxy, dimethylamino, methoxyethylamino, di-(methoxyethyl)amino or hydroxypropylamino,


and wherein any heterocyclyl group within a substituent on R¹ optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from methyl, ethyl, methoxy, ethoxy, formyl, acetyl, hydroxyl, hydroxymethyl, fluoroethyl or hydroxypropylamino,

and wherein any heterocyclyl group within a substituent on R¹ optionally bears 1 or 2 oxo substituents;


n is 0 or 1 and R³ group, if present, is located at the 5 position of the 1,3-benzodioxol group and is selected from fluoro or chloro;

10 Z² is a C≡C or CH=CH group; and

R¹⁴ is selected from (1-6C)alkoxy-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, hydroxyl-(1-6C)alkyl or from a group of formula:




15 wherein x is 0, 1, 2 or 3, R^c is hydrogen or (1-6C)alkyl and R^a and R^b are each independently selected from hydrogen and (1-6C)alkyl or R^a and R^b together with the nitrogen to which they are attached form morpholino, or from a group of the formula:



wherein X⁷ is a direct bond and Q⁵ is aryl, heteroaryl, heterocyclyl, heterocyclyl-(1-6C)alkyl, 20 or heterocycliloxy-(1-6C)alkyl,

and wherein any CH, CH₂ or CH₃ group within a R¹⁴ substituent optionally bears on each said CH, CH₂ or CH₃ group a substituent selected from (1-6C)alkoxy, carbamoyl, N-(1-6C)alkylcarbamoyl or a group of the formula:



25 wherein X⁸ is a direct bond or O and Q⁶ is (3-7C)cycloalkyl or heterocyclyl and wherein any heterocyclyl group within a substituent on R¹⁴ optionally bears 1 oxo substituent;

or a pharmaceutically acceptable acid addition salt thereof.

30 9. A quinazoline derivative of the Formula I according to claim 1 wherein Z is NH

- 164 -

m is 2 and the first **R**¹ group is at the 5-position and is selected from isopropoxy, tetrahydro-2H-pyran-4-yloxy and the second **R**¹ group is at the 7- position and is selected from

methoxy, 3-morpholin-4-ylpropoxy, 3-(4-acetylpiperazin-1-yl)propoxy, 3-(4-formylacetylpiperazin-1-yl)propoxy and 3-(3-oxo-4-methyl-piperazin-1-yl)propoxy

n is 1 and **R**³ group is located at the 5-position of the 1,3-benzodioxol-4-yl group and is chloro;

Z² is a C≡C or CH=CH group; and

R¹⁴ is selected from methoxymethyl, 2-methoxyethyl, methoxyisopropyl and pyridin-2-yl,

or a pharmaceutically acceptable acid addition salt thereof.

10. A quinazoline derivative of the Formula 1 according to claim 1 and selected from
 - N*-[5-chloro-7-(3-methoxyprop-1-ynyl)-1,3-benzodioxol-4-yl]-6-methoxy-7-(3-morpholin-4-ylpropoxy)quinazolin-4-amine,
 - N*-[5-chloro-7-(4-methoxybut-1-yn-1-yl)-1,3-benzodioxol-4-yl]-6-methoxy-7-(3-morpholin-4-ylpropoxy)quinazolin-4-amine,
 - N*-[5-chloro-7-(3-ethoxyprop-1-yn-1-yl)-1,3-benzodioxol-4-yl]-6-methoxy-7-(3-morpholin-4-ylpropoxy)quinazolin-4-amine,
 - N*-[5-chloro-7-(3-isopropoxyprop-1-yn-1-yl)-1,3-benzodioxol-4-yl]-6-methoxy-7-(3-morpholin-4-ylpropoxy)quinazolin-4-amine,
 - N*-[5-chloro-7-[3-(cyclopropylmethoxy)prop-1-yn-1-yl]-1,3-benzodioxol-4-yl]-6-methoxy-7-(3-morpholin-4-ylpropoxy)quinazolin-4-amine,
 - (1-{3-[(4-{[5-chloro-7-(4-methoxybut-1-yn-1-yl)-1,3-benzodioxol-4-yl]amino}-6-methoxyquinazolin-7-yl)oxy]propyl}piperidin-4-yl)methanol,
 - N'*-[3-(6-chloro-7-{[6-methoxy-7-(3-morpholin-4-ylpropoxy)quinazolin-4-yl]amino}-1,3-benzodioxol-4-yl)prop-2-yn-1-yl]-*N,N*-dimethylurea,
 - 7-{3-[bis(2-methoxyethyl)amino]propoxy}-*N*-[5-chloro-7-(4-methoxybut-1-yn-1-yl)-1,3-benzodioxol-4-yl]-6-methoxyquinazolin-4-amine,
 - 4-{3-[(4-{[5-chloro-7-(4-methoxybut-1-yn-1-yl)-1,3-benzodioxol-4-yl]amino}-6-methoxyquinazolin-7-yl)oxy]propyl}piperazine-1-carbaldehyde,

- N*-[5-chloro-7-(4-methoxybut-1-yn-1-yl)-1,3-benzodioxol-4-yl]-7-{3-[4-(2-fluoroethyl)piperazin-1-yl]propoxy}-6-methoxyquinazolin-4-amine,
- N*-[5-chloro-7-(4-methoxybut-1-yn-1-yl)-1,3-benzodioxol-4-yl]-6-methoxy-7-[3-(4-methoxypiperidin-1-yl)propoxy]quinazolin-4-amine,
- 5 4-{3-[(4-{[5-chloro-7-(4-methoxybut-1-yn-1-yl)-1,3-benzodioxol-4-yl]amino}-6-methoxyquinazolin-7-yl)oxy]propyl}-1-methylpiperazin-2-one,
- 4-{3-[(4-{[5-chloro-7-(3-methoxyprop-1-yn-1-yl)-1,3-benzodioxol-4-yl]amino}-6-methoxyquinazolin-7-yl)oxy]propyl}-1-methylpiperazin-2-one,
- N*-{3-[6-chloro-7-({6-methoxy-7-[3-(4-methyl-3-oxopiperazin-1-yl)propoxy]quinazolin-4-yl}amino)-1,3-benzodioxol-4-yl]prop-2-yn-1-yl}-*N,N*-dimethylurea,
- 10 1-{3-[(4-{[5-chloro-7-(pyridin-2-ylethynyl)-1,3-benzodioxol-4-yl]amino}-6-methoxyquinazolin-7-yl)oxy]propyl}-4-methylpiperazin-2-one,
- N*-{3-[6-chloro-7-({7-[3-(cis-2,6-dimethylmorpholin-4-yl)propoxy]-6-methoxyquinazolin-4-yl}amino)-1,3-benzodioxol-4-yl]prop-2-yn-1-yl}-*N,N*-dimethylurea,
- 15 *N*-[5-chloro-7-(3-methoxyprop-1-yn-1-yl)-1,3-benzodioxol-4-yl]-6-methoxy-7-(4-morpholin-4-ylbutoxy)quinazolin-4-amine,
- N*-[5-chloro-7-(pyridin-2-ylethynyl)-1,3-benzodioxol-4-yl]-6-methoxy-7-(3-morpholin-4-ylpropoxy)quinazolin-4-amine,
- N*-[5-chloro-7-(3-methoxyprop-1-yn-1-yl)-1,3-benzodioxol-4-yl]-7-{3-(cis-2,6-dimethylmorpholin-4-yl)propoxy}-6-methoxyquinazolin-4-amine,
- 20 *N*-[5-chloro-7-[(tetrahydrofuran-3-ylmethoxy)methyl]-1,3-benzodioxol-4-yl]-6-methoxy-7-(3-morpholin-4-ylpropoxy)quinazolin-4-amine,
- N*-[5-chloro-7-[(1,3-dioxolan-2-yl)methoxy)methyl]-1,3-benzodioxol-4-yl]-6-methoxy-7-(3-morpholin-4-ylpropoxy)quinazolin-4-amine,
- 25 *N*-[5-chloro-7-[(tetrahydrofuran-3-yloxy)methyl]-1,3-benzodioxol-4-yl]-6-methoxy-7-(3-morpholin-4-ylpropoxy)quinazolin-4-amine,
- N*-[5-chloro-7-(pyridin-3-ylethynyl)-1,3-benzodioxol-4-yl]-6-methoxy-7-(3-morpholin-4-ylpropoxy)quinazolin-4-amine,
- N*-[5-chloro-7-(1*H*-pyrazol-4-ylethynyl)-1,3-benzodioxol-4-yl]-6-methoxy-7-(3-morpholin-4-ylpropoxy)quinazolin-4-amine,
- 30

- 166 -

4-{3-[(4-{[5-chloro-7-(3-methoxybut-1-yn-1-yl)-1,3-benzodioxol-4-yl]amino}-6-methoxyquinazolin-7-yl)oxy]propyl}-1-methylpiperazin-2-one,

N-[5-bromo-7-(3-methoxybut-1-yn-1-yl)-1,3-benzodioxol-4-yl]-6-methoxy-7-(3-morpholin-4-ylpropoxy)quinazolin-4-amine,

5 4-{3-[(4-{[5-bromo-7-(3-methoxyprop-1-yn-1-yl)-1,3-benzodioxol-4-yl]amino}-6-methoxyquinazolin-7-yl)oxy]propyl}-1-methylpiperazin-2-one,

N-[5-chloro-7-(pyridin-2-ylethynyl)-1,3-benzodioxol-4-yl]-7-[3-(cis-2,6dimethylmorpholin-4-yl)propoxy]-6-methoxyquinazolin-4-amine,

4-{3-[(4-{[5-Chloro-7-(3-isopropoxyprop-1-yn-1-yl)-1,3-benzodioxol-4-yl]amino}-6-methoxyquinazolin-7-yl)oxy]propyl}-1-methylpiperazin-2-one,

10 *N*-[5-chloro-7-(1*H*-pyrazol-5-ylethynyl)-1,3-benzodioxol-4-yl]-6-methoxy-7-(3-morpholin-4-ylpropoxy)quinazolin-4-amine,

4-{3-[(4-{[5-chloro-7-(pyridin-2-ylethynyl)-1,3-benzodioxol-4-yl]amino}-6-methoxyquinazolin-7-yl)oxy]propyl}-1-methylpiperazin-2-one,

15 *N*-[5-chloro-7-(3-methoxyprop-1-yn-1-yl)-1,3-benzodioxol-4-yl]-6-methoxy-7-[3-(4-hydroxypiperidin-1-yl)propoxy]-quinazolin-4-amine,

((*2R*)-1-{3-[(4-{[5-chloro-7-(3-methoxyprop-1-yn-1-yl)-1,3-benzodioxol-4-yl]amino}-6-methoxyquinazolin-7-yl)oxy]propyl}pyrrolidin-2-yl)methanol,

4-{3-[(4-{[5-chloro-7-(3-methoxyprop-1-yn-1-yl)-1,3-benzodioxol-4-yl]amino}-6-methoxyquinazolin-7-yl)oxy]propyl}piperazin-2-one,

20 *N*-[5-chloro-7-(3-methoxyprop-1-yn-1-yl)-1,3-benzodioxol-4-yl]-7-[3-[4-(2-fluoroethyl)piperazin-1-yl]propoxy]-6-methoxyquinazolin-4-amine,

N-[5-chloro-7-(4-methoxybut-1-yn-1-yl)-1,3-benzodioxol-4-yl]-7-(3-morpholin-4-ylpropoxy)-5-(tetrahydro-2*H*-pyran-4-yloxy)quinazolin-4-amine,

25 *N*-[5-chloro-7-(4-methoxyprop-1-yn-1-yl)-1,3-benzodioxol-4-yl]-7-(3-morpholin-4-ylpropoxy)-5-isopropoxyquinazolin-4-amine,

7-[3-(4-acetylpiperazin-1-yl)propoxy]-*N*-[5-chloro-7-(3-methoxyprop-1-yn-1-yl)-1,3-benzodioxol-4-yl]-5-isopropoxyquinazolin-4-amine,

30 *N*-[5-chloro-7-(pyridin-2-ylethynyl)-1,3-benzodioxol-4-yl]-5-isopropoxy-7-(3-morpholin-4-ylpropoxy)quinazolin-4-amine,

- 167 -

4-{3-[(4-{[5-chloro-7-(3-methoxyprop-1-yn-1-yl)-1,3-benzodioxol-4-yl]amino}-5-isopropoxyquinazolin-7-yl)oxy]propyl}piperazine-1-carbaldehyde,

4-{3-[(4-{[5-chloro-7-(3-methoxyprop-1-yn-1-yl)-1,3-benzodioxol-4-yl]amino}-5-isopropoxyquinazolin-7-yl)oxy]propyl}-1-methylpiperazin-2-one,

5 *N*-[5-chloro-7-(3-methoxybut-1-yn-1-yl)-1,3-benzodioxol-4-yl]-6-methoxy-7-(3-morpholin-4-ylpropoxy)quinazolin-4-amine,


N-[5-bromo-7-(3-methoxyprop-1-yn-1-yl)-1,3-benzodioxol-4-yl]-6-methoxy-7-(3-morpholin-4-ylpropoxy)quinazolin-4-amine and

N-[5-chloro-7-(3-methoxyprop-1-yn-1-yl)-1,3-benzodioxol-4-yl]-7-[3-(dimethylamino)propoxy]-6-methoxyquinazolin-4-amine,

or a pharmaceutically acceptable acid addition salt thereof.


11. A process for the preparation of a quinazoline derivative of the Formula I or a pharmaceutically-acceptable salt thereof, according to claim 1 which comprises:-

15 (a) the reaction of a quinazoline of the Formula II



II

wherein L is a displaceable group and m and R¹ have any of the meanings defined in claim 1 except that any functional group is protected if necessary, with a compound of the Formula III



III

20


wherein Z is O, S, or N(R²) and n, R³, R², Z² and R¹⁴ have any of the meanings defined in claim 1 except that any functional group is protected if necessary, whereafter any protecting group that is present is removed by conventional means;

(b) for the production of those compounds of the Formula I wherein at least one R¹ group
25 is a group of the formula




- 168 -

wherein Q^1 is an aryl-(1-6C)alkyl, (3-7C)cycloalkyl-(1-6C)alkyl, (3-7C)cycloalkenyl-(1-6C)alkyl, heteroaryl-(1-6C)alkyl or heterocyclyl-(1-6C)alkyl group or an optionally substituted alkyl group and X^1 is an oxygen atom, the coupling, conveniently in the presence of a suitable dehydrating agent, of a quinazoline of the Formula V




- 5 wherein m , R^1 , Z , n , R^3 , Z^2 and R^{14} have any of the meanings defined in claim 1, except that any functional group is protected if necessary, with an appropriate alcohol of the formula Q^1-OH wherein any functional group is protected if necessary, whereafter any protecting group that is present is removed by conventional means;
- 10 (c) for the production of those compounds of the Formula I wherein R^1 is an amino-substituted (1-6C)alkoxy group (such as 2-homopiperidin-1-ylethoxy or 3-dimethylaminopropoxy), the reaction of a compound of the Formula I wherein R^1 is a halogeno-substituted (1-6C)alkoxy group with a heterocyclyl compound or an appropriate amine;
- 15 (d) for the production of those compounds of the Formula I wherein an R^1 group contains a (1-6C)alkoxy or substituted (1-6C)alkoxy group or a (1-6C)alkylamino or substituted (1-6C)alkylamino group, the alkylation of a quinazoline derivative of the Formula I wherein the R^1 group contains a hydroxy group or a primary or secondary amino group as appropriate;
- (e) for the production of those compounds of the Formula I wherein Z is a SO or SO_2 group, wherein an R^1 or R^3 substituent is a (1-6C)alkylsulphinyl or (1-6C)alkylsulphonyl group or wherein an R^1 , R^3 or R^{14} substituent contains a SO or SO_2 group, the oxidation of a compound of the Formula I wherein Z is a S group or wherein an R^1 or R^3 substituent is a (1-6C)alkylthio group or wherein an R^1 , R^3 or R^{14} substituent contains a S group as appropriate;
- 20 (f) the reaction of a compound of the Formula VI
- 25

- 169 -



VI

wherein L is a displaceable group as defined hereinbefore and m, R¹, Z, n and R³ have any of the meanings defined in claim hereinbefore except that any functional group is protected if necessary, with a compound of the Formula VII



VII


5

wherein Z² is a C≡C or C(R¹³)=C(R¹³) group and R¹³ and R¹⁴ have any of the meanings defined in claim 1 except that any functional group is protected if necessary, whereafter any protecting group that is present is removed by conventional means;

(g) for the production of a compound of the Formula I wherein R¹⁴ is a carboxy group, the cleavage of a compound of the Formula I wherein R¹⁴ is a (1-6C)alkoxycarbonyl group;

(h) the reaction of a compound of the Formula I wherein R¹⁴ is a carboxy group with an appropriate amine to form a further compound of the Formula I wherein R¹⁴ is a carbamoyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl or heterocyclylcarbonylamino group; or

15 (i) a coupling reaction of a compound of the Formula VIII



VIII

wherein m, R¹, Z, n and R³ have any of the meanings defined in claim 1 except that any functional group is protected if necessary, with a compound of the Formula IX



IX

- 170 -

wherein L is a displaceable group and R¹⁴ has any of the meanings defined in claim 1 except that any functional group is protected if necessary, whereafter any protecting group that is present is removed by conventional means;

and when a pharmaceutically-acceptable salt of a quinazoline derivative of Formula I
5 is required it may be obtained using a conventional procedure.

12. A pharmaceutical composition which comprises a quinazoline derivative of the Formula I, or a pharmaceutically-acceptable salt thereof as defined in claim 1 in association with a pharmaceutically-acceptable diluent or carrier.

10

13. A quinazoline derivative of the Formula I or a pharmaceutically acceptable salt thereof for use in a method of the treatment of the human or animal body by therapy.

14. A quinazoline derivative of the Formula I or a pharmaceutically acceptable salt
15 thereof, as defined in claim 1 for use in the treatment of cancer.

15. The use of a quinazoline derivative of the Formula I or a pharmaceutically acceptable salt thereof as defined in claim 1 in the manufacture of a medicament for use as an anti-proliferative agent in the containment and/or treatment of solid tumour disease.

20

16. The use of a quinazoline derivative of the Formula I or a pharmaceutically acceptable salt thereof as defined in claim 1 in the manufacture of a medicament for use as an anti-invasive agent in the containment and/or treatment of solid tumour disease.

25